

TECHNICAL INFORMATION BULLETIN

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MINNESOTA, MISSISSIPPI, MISSOURI, NEBRASKA, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, VIRGINIA, WEST VIRGINIA and WISCONSIN.

BALLAD[®] PLUS **EPA Reg. No. 69592-13**

For use of Ballad[®] Plus on soybeans at reduced application rates for control of the diseases specified in the accompanying application table.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This recommendation must be in the possession of the user at the time of pesticide application.

To enhance plant disease control, reduced rates of Ballad Plus alone may be applied to soybeans. Ballad Plus cannot be mixed with any product with prohibition against such mixing. When tank-mixing Ballad Plus with other registered pesticides, always read and follow all use directions, restrictions, and precautions of both Ballad Plus and the tank-mix partner(s). Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. Do not exceed label dosage rates for the tank mix partner and follow the dosage rates for Ballad Plus set forth in this recommendation.

Application Rates of Ballad Plus for Use on Soybeans			
Crops	Diseases	Rate qts./A	Application Instructions
Oil Seed Crops Soybeans	Bacterial Speck <i>Pseudomonas syringae</i> pv. <i>glycinea</i> Brown Spot <i>Septoria glycines</i> Cercospora Leaf Spot <i>Cercospora</i> spp. Pod and Stem Blight <i>Diaporthe phaseolorum</i> var. <i>sojæ</i> <i>Phomopsis longicola</i> Downy Mildew <i>Peronospora manshurica</i> White Mold (Sclerotinia Stem Rot) <i>Sclerotinia sclerotiorum</i> Bacterial Pustule <i>Xanthomonas</i> spp. Asian Soybean Rust <i>Phakopsora pachyrhizi</i>	0.5 - 4	Begin applications of Ballad Plus when conditions are conducive to disease development. Repeat on 7 to 14 day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure.

Ballad® is a registered trademark of AgraQuest, Inc.
 © Copyright AgraQuest, Inc., 2011

072011

AgraQuest, Inc.
 1540 Drew Avenue
 Davis, California 95618
 (530) 750-0150
 www.agraquest.com

